



INSTRUCTIONS FOR THE iKnowMed® ELECTRONIC HEALTH RECORD (EHR) SYSTEM

UPDATING RELAPSED OR REFRACTORY MULTIPLE MYELOMA ORDER SETS WITH TALVEY™ (talquetamab-tgvs)

INDICATION AND USAGE

TALVEY™ (talquetamab-tgvs) is indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least four prior lines of therapy, including a proteasome inhibitor, an immunomodulatory agent, and an anti-CD38 monoclonal antibody.

This indication is approved under accelerated approval based on response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

SELECT IMPORTANT SAFETY INFORMATION

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGIC TOXICITY, including IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

Cytokine release syndrome (CRS), including life-threatening or fatal reactions, can occur in patients receiving TALVEY™. Initiate TALVEY™ treatment with step-up dosing to reduce the risk of CRS. Withhold TALVEY™ until CRS resolves or permanently discontinue based on severity.

Neurologic toxicity, including immune effector cell-associated neurotoxicity syndrome (ICANS), and serious and life-threatening or fatal reactions, can occur with TALVEY™. Monitor patients for signs and symptoms of neurologic toxicity including ICANS during treatment. Withhold or discontinue TALVEY™ based on severity.

Because of the risk of CRS and neurologic toxicity, including ICANS, TALVEY™ is available only through a restricted program called the TECVAYLI® and TALVEY™ Risk Evaluation and Mitigation Strategy (REMS).

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.

CONTENTS

1. Overview and Limitations	3
2. Background and Indication	3
3. Considerations	4
Important Dosing Information	4
Recommended Dosage	5
Recommended Pretreatment Medications	6
Dosage Modifications for Adverse Reactions	6
4. iKnowMed® Electronic Health Record Instructions	7
5. Disclaimers	20
6. Important Safety Information	21
7. Notes	24

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



1. Overview and Limitations

This document is intended to provide health systems with instructions to update relapsed or refractory multiple myeloma order sets with TALVEY™ (talquetamab-tgvs), within the approved Indication and consistent with the Prescribing Information. This document is not intended to provide any clinical advice or clinical recommendations, which are solely the responsibility of the health system.

These instructions are specific to adult relapsed or refractory multiple myeloma and to the iKnowMed® EHR system, and are not appropriate for other conditions, treatments, therapeutic areas, or for other EHR systems.

The process outlined on the following pages is variable and not all steps will apply to every health system. Any steps or settings outlined in this document that are not part of a health system's standard process should be excluded or modified accordingly. Any questions should be directed to the appropriate service provider. The practice is solely responsible for implementing, testing, monitoring, and the ongoing operation of any EHR tools.

2. Background and Indication

Janssen Biotech, Inc., has developed this resource for health systems who desire to update EHR order sets with TALVEY™.

INDICATION AND USAGE

TALVEY™ (talquetamab-tgvs) is indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least four prior lines of therapy, including a proteasome inhibitor, an immunomodulatory agent, and an anti-CD38 monoclonal antibody.

This indication is approved under accelerated approval based on response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



3. Considerations

The clinical data elements provided are only suggestions, and it is strongly recommended that clinical and operational leadership determine that the final elements align with the expectations and goals of the organization. A Notes section (located on [page 24](#) of this guide) can be used by clinical leadership to document any additions or changes that need to be made in order to align these instructions with practice protocols or pathways.

Consider reviewing appropriate order sets for the inclusion of TALVEY™. Order sets may benefit from regular updates to include new treatments, laboratory sets, and other orderable items.

End users of the order sets should be offered an update of the contents and availability of the new order sets (and training if needed). In many cases, end users will have already been trained on how to use order sets as part of the health system's best practices, thereby reducing the need for additional training.

The following provides an overview and quick reference guide for the dosing and administration details for TALVEY™ for its approved indication as per the Prescribing Information. Step-by-step instructions for applying these medication details in the order set update are shown in the following pages of this guide.

Important Dosing Information

Administer TALVEY™ subcutaneously according to the step-up dosing schedule in Tables 1 and 2 to reduce the incidence and severity of cytokine release syndrome (CRS).

Administer pretreatment medications prior to each dose of TALVEY™ in the step-up dosing schedule as recommended.

TALVEY™ should only be administered by a qualified healthcare professional with appropriate medical support to manage severe reactions such as CRS and neurologic toxicity including immune effector cell-associated neurotoxicity syndrome (ICANS).

Due to the risk of CRS and neurologic toxicity, including ICANS, patients should be hospitalized for 48 hours after administration of all doses within the TALVEY™ step-up dosing schedule.

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



Recommended Dosage

For subcutaneous injection.

Administer pretreatment medications prior to each dose of TALVEY™ in the step-up dosing schedule.

Administer TALVEY™ subcutaneously on a weekly or biweekly (every 2 weeks) dosing schedule according to Table 1 or Table 2 in the full Prescribing Information. Continue treatment until disease progression or unacceptable toxicity.

Table 1: TALVEY™ Weekly Dosing Schedule

Dosing schedule	Day	Dose ^a	
Step-up dosing schedule	Day 1	Step-up dose 1	0.01 mg/kg
	Day 4 ^b	Step-up dose 2	0.06 mg/kg
	Day 7 ^b	First treatment dose	0.4 mg/kg
Weekly dosing schedule	One week after first treatment dose and weekly thereafter ^c	Subsequent treatment doses	0.4 mg/kg once weekly

^aBased on actual body weight.

^bDose may be administered between 2 to 4 days after the previous dose and may be given up to 7 days after the previous dose to allow for resolution of adverse reactions.

^cMaintain a minimum of 6 days between weekly doses.

Table 2: TALVEY™ Biweekly (Every 2 Weeks) Dosing Schedule

Dosing schedule	Day	Dose ^a	
Step-up dosing schedule	Day 1	Step-up dose 1	0.01 mg/kg
	Day 4 ^b	Step-up dose 2	0.06 mg/kg
	Day 7 ^b	Step-up dose 3	0.4 mg/kg
	Day 10 ^c	First treatment dose	0.8 mg/kg
Biweekly (every 2 weeks) dosing schedule	Two weeks after first treatment dose and every 2 weeks thereafter ^d	Subsequent treatment doses	0.8 mg/kg every 2 weeks

^aBased on actual body weight.

^bDose may be administered between 2 to 4 days after the previous dose and may be given up to 7 days after the previous dose to allow for resolution of adverse reactions.

^cDose may be administered between 2 to 7 days after step-up dose 3.

^dMaintain a minimum of 12 days between biweekly (every 2 weeks) doses.

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



Recommended Pretreatment Medications

Administer the following pretreatment medications 1 to 3 hours before each dose of TALVEY™ in the step-up dosing schedule to reduce the risk of CRS:

- Corticosteroid (oral or intravenous dexamethasone, 16 mg or equivalent)
- Antihistamines (oral or intravenous diphenhydramine, 50 mg or equivalent)
- Antipyretics (oral or intravenous acetaminophen, 650 mg to 1,000 mg or equivalent)

Administration of pretreatment medications may be required for subsequent doses of TALVEY™ in the following patients:

- Patients who repeat doses within the TALVEY™ step-up dosing schedule due to dose delays (see Table 3 or Table 4 in the full Prescribing Information)
- Patients who experienced CRS (see Table 5 in the full Prescribing Information)

Dosage Modifications for Adverse Reactions

Dose delays may be required to manage toxicities related to TALVEY™.

See Dosage and Administration Section 2.4 in the full Prescribing Information for guidance regarding restarting TALVEY™ after dosage delay.

See Table 5, Table 6, and Table 7 in the full Prescribing Information for recommended actions for the management of CRS, ICANS, and neurologic toxicity. See Table 8 in the full Prescribing Information for recommended dose modifications for other adverse reactions.

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



4. iKnowMed® Electronic Health Record Instructions

Treatment Regimens are commonly used in the management of oncology patients. After initial release, Treatment Regimens may benefit from a clinical update. The optimization of Treatment Regimens is a common process and provides an opportunity to incorporate treatment updates.

The iKnowMed EHR allows for customizing a Treatment Regimen 1) outside of Clear Value Plus or 2) using the Clear Value Plus exception. Customers who do not use Clear Value Plus should proceed with the instructions in section 4.1 starting below. Customers who do use Clear Value Plus should proceed with the instructions in section 4.2 starting on [page 13](#).

There are 2 options to optimize a Treatment Regimen in the iKnowMed EHR system.

4.1. Optimizing a Treatment Regimen for Customers Not Using Clear Value Plus

4.2. Optimizing a Treatment Regimen for Customers Using the Exception in Clear Value Plus

4.1 Optimizing a Treatment Regimen for Customers Not Using Clear Value Plus

4.1.1. Access the Regimens Templates with admin credentials (Treatment Regimen Orders editor).

4.1.2. In the Regimen Search window, **Search by Diagnosis/Problem** by entering **“relapsed or refractory multiple myeloma”** in the Search field. **Click Search for Regimens** to display Treatment Regimens matching the search terms. Select the desired Treatment Regimen to optimize by copying and renaming it.

4.1.3. Adjust the Regimens Search Rules and Problems Associations if desired.

4.1.4. Complete the Display Name as “TALVEY™ for relapsed or refractory multiple myeloma.”

4.1.5. Add in the Reference Name as **“TALVEY™ (talquetamab-tgvs) is indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least four prior lines of therapy, including a proteasome inhibitor, an immunomodulatory agent, and an anti-CD38 monoclonal antibody.”**

→ *See next page for additional steps.*

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



4.1.6. In the Regimen Comments or Instructions to Provider field, **enter:**

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGIC TOXICITY, including IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

Cytokine release syndrome (CRS), including life-threatening or fatal reactions, can occur in patients receiving TALVEY™. Initiate TALVEY™ treatment with step-up dosing to reduce the risk of CRS. Withhold TALVEY™ until CRS resolves or permanently discontinue based on severity.

Neurologic toxicity, including immune effector cell-associated neurotoxicity syndrome (ICANS), and serious and life-threatening or fatal reactions, can occur with TALVEY™. Monitor patients for signs and symptoms of neurologic toxicity including ICANS during treatment. Withhold or discontinue TALVEY™ based on severity.

Because of the risk of CRS and neurologic toxicity, including ICANS, TALVEY™ is available only through a restricted program called the TECVAYLI® and TALVEY™ Risk Evaluation and Mitigation Strategy (REMS).

Important Dosing Information:

- Administer TALVEY™ subcutaneously according to the step-up dosing schedule in Tables 1 and 2 to reduce the incidence and severity of cytokine release syndrome (CRS)
- Administer pretreatment medications prior to each dose of TALVEY™ in the step-up dosing schedule as recommended
- TALVEY™ should only be administered by a qualified healthcare professional with appropriate medical support to manage severe reactions such as CRS and neurologic toxicity including immune effector cell-associated neurotoxicity syndrome (ICANS)
- Due to the risk of CRS and neurologic toxicity, including ICANS, patients should be hospitalized for 48 hours after administration of all doses within the TALVEY™ step-up dosing schedule

Recommended Dosage:

- For subcutaneous injection
- Administer pretreatment medications prior to each dose of TALVEY™ in the step-up dosing schedule
- Administer TALVEY™ subcutaneously on a weekly or biweekly (every 2 weeks) dosing schedule according to Table 1 or Table 2 in the full Prescribing Information. Continue treatment until disease progression or unacceptable toxicity

→ See next page for additional steps.

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



4.1.6. (continued)

Table 1: TALVEY™ Weekly Dosing Schedule

Dosing schedule	Day	Dose ^a	
Step-up dosing schedule	Day 1	Step-up dose 1	0.01 mg/kg
	Day 4 ^b	Step-up dose 2	0.06 mg/kg
	Day 7 ^b	First treatment dose	0.4 mg/kg
Weekly dosing schedule	One week after first treatment dose and weekly thereafter ^c	Subsequent treatment doses	0.4 mg/kg once weekly

^aBased on actual body weight.

^bDose may be administered between 2 to 4 days after the previous dose and may be given up to 7 days after the previous dose to allow for resolution of adverse reactions.

^cMaintain a minimum of 6 days between weekly doses.

Table 2: TALVEY™ Biweekly (Every 2 Weeks) Dosing Schedule

Dosing schedule	Day	Dose ^a	
Step-up dosing schedule	Day 1	Step-up dose 1	0.01 mg/kg
	Day 4 ^b	Step-up dose 2	0.06 mg/kg
	Day 7 ^b	Step-up dose 3	0.4 mg/kg
	Day 10 ^c	First treatment dose	0.8 mg/kg
Biweekly (every 2 weeks) dosing schedule	Two weeks after first treatment dose and every 2 weeks thereafter ^d	Subsequent treatment doses	0.8 mg/kg every 2 weeks

^aBased on actual body weight.

^bDose may be administered between 2 to 4 days after the previous dose and may be given up to 7 days after the previous dose to allow for resolution of adverse reactions.

^cDose may be administered between 2 to 7 days after step-up dose 3.

^dMaintain a minimum of 12 days between biweekly (every 2 weeks) doses.

Dosage Modifications for Adverse Reactions:

- Dose delays may be required to manage toxicities related to TALVEY™
- See Dosage and Administration Section 2.4 in the full Prescribing Information for guidance regarding restarting TALVEY™ after dosage delay
- See Table 5, Table 6, and Table 7 in the full Prescribing Information for recommended actions for the management of CRS, ICANS, and neurologic toxicity. See Table 8 in the full Prescribing Information for recommended dose modifications for other adverse reactions

→ See next page for additional steps.

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



4.1.7. Click + Add Group and add a Treatment Group for the TALVEY™ Premedications:

Administer the following pretreatment medications 1 to 3 hours before each dose of TALVEY™ in the step-up dosing schedule to reduce the risk of CRS:

- Corticosteroid (oral or intravenous dexamethasone, 16 mg or equivalent)
- Antihistamines (oral or intravenous diphenhydramine, 50 mg or equivalent)
- Antipyretics (oral or intravenous acetaminophen, 650 mg to 1,000 mg or equivalent)
- ◆ Complete the medication details (SIG, dose, route, frequency, offset time, admin over, and free form any special Admin instructions to the nurse)
- ◆ Administration of pretreatment medications may be required for subsequent doses for patients who repeat doses within the TALVEY™ step-up dosing schedule due to dose delays (see Table 3 or Table 4 in the full Prescribing Information) or for patients who experienced CRS (see Table 5 in the full Prescribing Information)

4.1.8. Add TALVEY™ to the desired Treatment Group by clicking the plus sign next to the Treatment Group.

4.1.9. In the **References**, enter:

For important information about administering TALVEY™, refer to the full Prescribing Information (<https://www.janssenlabels.com/package-insert/product-monograph/prescribing-information/TALVEY-pi.pdf>).

TALVEY™ REMS: TALVEY™ is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called the TECVAYLI® and TALVEY™ REMS because of the risks of CRS and neurologic toxicity, including ICANS.

Notable requirements of the TECVAYLI® and TALVEY™ REMS include the following:

- Prescribers must be certified with the program by enrolling and completing training
- Prescribers must counsel patients receiving TALVEY™ about the risk of CRS and neurologic toxicity, including ICANS and provide patients with Patient Wallet Card
- Pharmacies and healthcare settings that dispense TALVEY™ must be certified with the TECVAYLI® and TALVEY™ REMS program and must verify prescribers are certified through the TECVAYLI® and TALVEY™ REMS program
- Wholesalers and distributors must only distribute TALVEY™ to certified pharmacies

Further information about the TECVAYLI® and TALVEY™ REMS program is available at www.TEC-TALREMS.com or by telephone at 1-855-810-8064.

→ See next page for additional steps.

Please read full Important Safety Information on **pages 21-23** and full **Prescribing Information**, including **Boxed WARNING**, for TALVEY™.



4.1.9. (continued)

Janssen Care Navigator Program: Janssen has resources to help you support your patients, including personalized 1-on-1 support.

Janssen Care Navigators can help patients:

- Discover cost support options that may help them pay for their medication
- Learn more about their cancer and Janssen medication
- Connect them to resources for their practical and emotional support

Schedule your patient for an introductory call from a Care Navigator here:

JanssenCompass.com/signup or make a referral over the phone by calling 1-844-628-1234.

Transitions of Care: Consider planning for patients' smooth transition between step-up dosing and weekly or biweekly (every 2 weeks) dosing, including necessary coverage and reimbursement planning.

4.1.10. In the Order Review tab, **complete the Treatment Regimen** and **add**:

– Weekly dosing schedule

- ◆ The recommended dosage of TALVEY™ (based on actual body weight) is step-up doses of 0.01 mg/kg, 0.06 mg/kg, and 0.4 mg/kg, followed by a weekly dose of 0.4 mg/kg with treatment continued once weekly until disease progression or unacceptable toxicity. See Table 3 in the full Prescribing Information for recommendations on restarting TALVEY™ after dose delays for adverse reactions
- ◆ Step-up dosing
 - Day 1: Step-up dose 1
 - » TALVEY™: 0.01 mg/kg
 - Day 4: Step-up dose 2
 - » TALVEY™: 0.06 mg/kg—step-up dose 2 may be administered between 2 to 4 days after step-up dose 1 and may be given up to 7 days after step-up dose 1 to allow for resolution of adverse reactions
 - Day 7: First treatment dose
 - » TALVEY™: 0.4 mg/kg—first treatment dose may be administered between 2 to 4 days after step-up dose 2 and may be given up to 7 days after step-up dose 2 to allow for resolution of adverse reactions
- ◆ Weekly dosing
 - One week after first treatment dose and weekly thereafter: Subsequent treatment doses
 - » TALVEY™: 0.4 mg/kg once weekly—maintain a minimum of 6 days between weekly doses

→ See next page for additional steps.

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



4.1.10. (continued)

– Biweekly (every 2 weeks) dosing schedule

- ◆ The recommended dosage of TALVEY™ (based on actual body weight) is step-up doses of 0.01 mg/kg, 0.06 mg/kg, 0.4 mg/kg, and 0.8 mg/kg, followed by a biweekly dose of 0.8 mg/kg with treatment continued once every two weeks until disease progression or unacceptable toxicity. See Table 4 in the full Prescribing Information for recommendations on restarting TALVEY™ after dose delays for adverse reactions
- ◆ Step-up dosing
 - Day 1: Step-up dose 1
 - » TALVEY™: 0.01 mg/kg
 - Day 4: Step-up dose 2
 - » TALVEY™: 0.06 mg/kg—step-up dose 2 may be administered between 2 to 4 days after step-up dose 1 and may be given up to 7 days after step-up dose 1 to allow for resolution of adverse reactions
 - Day 7: Step-up dose 3
 - » TALVEY™: 0.4 mg/kg—step-up dose 3 may be administered between 2 to 4 days after step-up dose 2 and may be given up to 7 days after step-up dose 2 to allow for resolution of adverse reactions
 - Day 10: First treatment dose
 - » TALVEY™: 0.8 mg/kg—first treatment dose may be administered between 2 to 7 days after step-up dose 3
- ◆ Biweekly (every 2 weeks) dosing
 - Two weeks after first treatment dose and every 2 weeks thereafter: Subsequent treatment doses
 - » TALVEY™: 0.8 mg/kg every 2 weeks—maintain a minimum of 12 days between biweekly (every 2 weeks) doses

4.1.11. Click **Save** to complete the new Treatment Regimen.

→ See next page for additional steps.

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including **Boxed WARNING**, for TALVEY™.



4.2 Optimizing a Treatment Regimen for Customers Using the Exception in Clear Value Plus

- 4.2.1. **Open a patient chart and select Order Regimen** in the Value Pathways section.
- 4.2.2. **Enter “relapsed or refractory multiple myeloma”** in the Search field. **Click Search** to display Treatment Regimens matching the search terms. Select the desired Treatment Regimen.
- 4.2.3. **Select Chemotherapy/Immunotherapy/Targeted Therapy** from the Regimen Type menu.
- 4.2.4. **Complete the Clear Value Plus wizard** to finalize the pathway information. On the left-hand side, a summary will build while progressing through the wizard. The Treatment Options field will list all available options.
- 4.2.5. **Select Record Warranted Exception** for an off-pathway treatment.
- 4.2.6. **Select the desired Exception Reason** and **enter a reason** (for example, “new research available”).
- 4.2.7. **Click Submit Warranted Exception.**
- 4.2.8. **Click Add New** in the Treatment Regimen header. Find TALVEY™ and **add it to the Treatment Regimen.**
- 4.2.9. **Complete the TALVEY™ medication details** (SIG, dose, route, frequency, offset time, admin over, and free form any special Admin instructions for the nurse). In the Instructions field, add a link to the TALVEY™ full Prescribing Information (<https://www.janssenlabels.com/package-insert/product-monograph/prescribing-information/TALVEY-pi.pdf>).

→ See next page for additional steps.

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



4.2.10. In the Regimen Comments or Instructions to Provider field, **enter**:

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGIC TOXICITY, including IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

Cytokine release syndrome (CRS), including life-threatening or fatal reactions, can occur in patients receiving TALVEY™. Initiate TALVEY™ treatment with step-up dosing to reduce the risk of CRS. Withhold TALVEY™ until CRS resolves or permanently discontinue based on severity.

Neurologic toxicity, including immune effector cell-associated neurotoxicity syndrome (ICANS), and serious and life-threatening or fatal reactions, can occur with TALVEY™. Monitor patients for signs and symptoms of neurologic toxicity including ICANS during treatment. Withhold or discontinue TALVEY™ based on severity.

Because of the risk of CRS and neurologic toxicity, including ICANS, TALVEY™ is available only through a restricted program called the TECVAYLI® and TALVEY™ Risk Evaluation and Mitigation Strategy (REMS).

Important Dosing Information:

- Administer TALVEY™ subcutaneously according to the step-up dosing schedule in Tables 1 and 2 to reduce the incidence and severity of cytokine release syndrome (CRS)
- Administer pretreatment medications prior to each dose of TALVEY™ in the step-up dosing schedule as recommended
- TALVEY™ should only be administered by a qualified healthcare professional with appropriate medical support to manage severe reactions such as CRS and neurologic toxicity including immune effector cell-associated neurotoxicity syndrome (ICANS)
- Due to the risk of CRS and neurologic toxicity, including ICANS, patients should be hospitalized for 48 hours after administration of all doses within the TALVEY™ step-up dosing schedule

Recommended Dosage:

- For subcutaneous injection
- Administer pretreatment medications prior to each dose of TALVEY™ in the step-up dosing schedule
- Administer TALVEY™ subcutaneously on a weekly or biweekly (every 2 weeks) dosing schedule according to Table 1 in the full Prescribing Information. Continue treatment until disease progression or unacceptable toxicity

→ See next page for additional steps.

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



4.2.10. (continued)

Table 1: TALVEY™ Weekly Dosing Schedule

Dosing schedule	Day	Dose ^a	
Step-up dosing schedule	Day 1	Step-up dose 1	0.01 mg/kg
	Day 4 ^b	Step-up dose 2	0.06 mg/kg
	Day 7 ^b	First treatment dose	0.4 mg/kg
Weekly dosing schedule	One week after first treatment dose and weekly thereafter ^c	Subsequent treatment doses	0.4 mg/kg once weekly

^aBased on actual body weight.

^bDose may be administered between 2 to 4 days after the previous dose and may be given up to 7 days after the previous dose to allow for resolution of adverse reactions.

^cMaintain a minimum of 6 days between weekly doses.

Table 2: TALVEY™ Biweekly (Every 2 Weeks) Dosing Schedule

Dosing schedule	Day	Dose ^a	
Step-up dosing schedule	Day 1	Step-up dose 1	0.01 mg/kg
	Day 4 ^b	Step-up dose 2	0.06 mg/kg
	Day 7 ^b	Step-up dose 3	0.4 mg/kg
	Day 10 ^c	First treatment dose	0.8 mg/kg
Biweekly (every 2 weeks) dosing schedule	Two weeks after first treatment dose and every 2 weeks thereafter ^d	Subsequent treatment doses	0.8 mg/kg every 2 weeks

^aBased on actual body weight.

^bDose may be administered between 2 to 4 days after the previous dose and may be given up to 7 days after the previous dose to allow for resolution of adverse reactions.

^cDose may be administered between 2 to 7 days after step-up dose 3.

^dMaintain a minimum of 12 days between biweekly (every 2 weeks) doses.

Dosage Modifications for Adverse Reactions:

- Dose delays may be required to manage toxicities related to TALVEY™
- See Dosage and Administration Section 2.4 in the full Prescribing Information for guidance regarding restarting TALVEY™ after dosage delay
- See Table 5, Table 6, and Table 7 in the full Prescribing Information for recommended actions for the management of CRS, ICANS, and neurologic toxicity. See Table 8 in the full Prescribing Information for recommended dose modifications for other adverse reactions

→ See next page for additional steps.

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



4.2.11. In the Schedule column, **click TALVEY™** to edit the Treatment Regimen Order:

– Weekly dosing schedule

- ◆ The recommended dosage of TALVEY™ (based on actual body weight) is step-up doses of 0.01 mg/kg, 0.06 mg/kg, and 0.4 mg/kg, followed by a weekly dose of 0.4 mg/kg with treatment continued once weekly until disease progression or unacceptable toxicity. See Table 3 in the full Prescribing Information for recommendations on restarting TALVEY™ after dose delays for adverse reactions
- ◆ Step-up dosing
 - Day 1: Step-up dose 1
 - » TALVEY™: 0.01 mg/kg
 - Day 4: Step-up dose 2
 - » TALVEY™: 0.06 mg/kg—step-up dose 2 may be administered between 2 to 4 days after step-up dose 1 and may be given up to 7 days after step-up dose 1 to allow for resolution of adverse reactions
 - Day 7: First treatment dose
 - » TALVEY™: 0.4 mg/kg—first treatment dose may be administered between 2 to 4 days after step-up dose 2 and may be given up to 7 days after step-up dose 2 to allow for resolution of adverse reactions
- ◆ Weekly dosing
 - One week after first treatment dose and weekly thereafter: Subsequent treatment doses
 - » TALVEY™: 0.4 mg/kg once weekly—maintain a minimum of 6 days between weekly doses

→ See next page for additional steps.

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



4.2.11. (continued)

– Biweekly (every 2 weeks) dosing schedule

- ◆ The recommended dosage of TALVEY™ (based on actual body weight) is step-up doses of 0.01 mg/kg, 0.06 mg/kg, 0.4 mg/kg, and 0.8 mg/kg, followed by a biweekly dose of 0.8 mg/kg with treatment continued once every two weeks until disease progression or unacceptable toxicity. See Table 4 in the full Prescribing Information for recommendations on restarting TALVEY™ after dose delays for adverse reactions
- ◆ Step-up dosing
 - Day 1: Step-up dose 1
 - » TALVEY™: 0.01 mg/kg
 - Day 4: Step-up dose 2
 - » TALVEY™: 0.06 mg/kg—step-up dose 2 may be administered between 2 to 4 days after step-up dose 1 and may be given up to 7 days after step-up dose 1 to allow for resolution of adverse reactions
 - Day 7: Step-up dose 3
 - » TALVEY™: 0.4 mg/kg—step-up dose 3 may be administered between 2 to 4 days after step-up dose 2 and may be given up to 7 days after step-up dose 2 to allow for resolution of adverse reactions
 - Day 10: First treatment dose
 - » TALVEY™: 0.8 mg/kg—first treatment dose may be administered between 2 to 7 days after step-up dose 3
- ◆ Biweekly (every 2 weeks) dosing
 - Two weeks after first treatment dose and every 2 weeks thereafter: Subsequent treatment doses
 - » TALVEY™: 0.8 mg/kg every 2 weeks—maintain a minimum of 12 days between biweekly (every 2 weeks) doses

→ See next page for additional steps.

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



4.2.12. In the **reference text field**, enter:

For important information about administering TALVEY™, refer to the full Prescribing Information (<https://www.janssenlabels.com/package-insert/product-monograph/prescribing-information/TALVEY-pi.pdf>).

TALVEY™ REMS: TALVEY™ is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called the TECVAYLI® and TALVEY™ REMS because of the risks of CRS and neurologic toxicity, including ICANS.

Notable requirements of the TECVAYLI® and TALVEY™ REMS include the following:

- Prescribers must be certified with the program by enrolling and completing training
- Prescribers must counsel patients receiving TALVEY™ about the risk of CRS and neurologic toxicity, including ICANS and provide patients with Patient Wallet Card
- Pharmacies and healthcare settings that dispense TALVEY™ must be certified with the TECVAYLI® and TALVEY™ REMS program and must verify prescribers are certified through the TECVAYLI® and TALVEY™ REMS program
- Wholesalers and distributors must only distribute TALVEY™ to certified pharmacies

Further information about the TECVAYLI® and TALVEY™ REMS program is available at www.TEC-TALREMS.com or by telephone at 1-855-810-8064.

Janssen Care Navigator Program: Janssen has resources to help you support your patients, including personalized 1-on-1 support.

Janssen Care Navigators can help patients:

- Discover cost support options that may help them pay for their medication
- Learn more about their cancer and Janssen medication
- Connect them to resources for their practical and emotional support

Schedule your patient for an introductory call from a Care Navigator here: JanssenCompass.com/signup or make a referral over the phone by calling 1-844-628-1234.

Transitions of Care: Consider planning for patients' smooth transition between step-up dosing and weekly or biweekly (every 2 weeks) dosing, including necessary coverage and reimbursement planning.

→ See next page for additional steps.

Please read full Important Safety Information on **pages 21-23** and full **Prescribing Information**, including **Boxed WARNING**, for TALVEY™.



4.2.13. Add the TALVEY™ Premedications:

Administer the following pretreatment medications 1 to 3 hours before each dose of TALVEY™ in the step-up dosing schedule to reduce the risk of CRS:

- Corticosteroid (oral or intravenous dexamethasone, 16 mg or equivalent)
- Antihistamines (oral or intravenous diphenhydramine, 50 mg or equivalent)
- Antipyretics (oral or intravenous acetaminophen, 650 mg to 1,000 mg or equivalent)
- ◆ Complete the medication details (SIG, dose, route, frequency, offset time, admin over, and free form any special Admin instructions to the nurse)
- ◆ Administration of pretreatment medications may be required for subsequent doses for patients who repeat doses within the TALVEY™ step-up dosing schedule due to dose delays (see Table 3 or Table 4 in the full Prescribing Information) or for patients who experienced CRS (see Table 5 in the full Prescribing Information)

4.2.14. Rename the new Treatment Regimen as “TALVEY™ (talquetamab-tgvs) is indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least four prior lines of therapy, including a proteasome inhibitor, an immunomodulatory agent, and an anti-CD38 monoclonal antibody” and click Close.

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



5. Disclaimers

- The Customer (eg, the medical group, Integrated Delivery Network [IDN], Organized Customer Group, and/or Health System) shall be solely responsible for implementation, testing, and monitoring of the instructions to ensure proper orientation in each Customer's EHR system
- Capabilities, functionality, and set-up (customization) for each individual EHR system vary. Janssen Biotech, Inc., shall not be responsible for revising the implementation instructions it provides to any Customer in the event that the Customer modifies or changes its software, or the configuration of its EHR system, after such time as the implementation instructions have been initially provided by Janssen Biotech, Inc. The instructions are not guaranteed to work for all available EHR systems, and Janssen Biotech, Inc., shall have no liability thereto
- While EHRs may assist providers in identifying appropriate patients for consideration of assessment and treatment, the decision and action should ultimately be decided by a provider in consultation with the patient, after a review of the patient's records to determine eligibility, and Janssen Biotech, Inc., shall have no liability thereto
- All products are trademarks of their respective holders, all rights reserved. Reference to these products is not intended to imply affiliation with or sponsorship of Janssen Biotech, Inc., and/or its affiliates

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.



6. Important Safety Information

INDICATION AND USAGE

TALVEY™ (talquetamab-tgvs) is indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least four prior lines of therapy, including a proteasome inhibitor, an immunomodulatory agent, and an anti-CD38 monoclonal antibody.

This indication is approved under accelerated approval based on response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

IMPORTANT SAFETY INFORMATION

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGIC TOXICITY, including IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

Cytokine release syndrome (CRS), including life-threatening or fatal reactions, can occur in patients receiving TALVEY™. Initiate TALVEY™ treatment with step-up dosing to reduce the risk of CRS. Withhold TALVEY™ until CRS resolves or permanently discontinue based on severity.

Neurologic toxicity, including immune effector cell-associated neurotoxicity syndrome (ICANS), and serious and life-threatening or fatal reactions, can occur with TALVEY™. Monitor patients for signs and symptoms of neurologic toxicity including ICANS during treatment. Withhold or discontinue TALVEY™ based on severity.

Because of the risk of CRS and neurologic toxicity, including ICANS, TALVEY™ is available only through a restricted program called the TECVAYLI® and TALVEY™ Risk Evaluation and Mitigation Strategy (REMS).

CONTRAINDICATIONS: None.

WARNINGS AND PRECAUTIONS

Cytokine Release Syndrome (CRS): TALVEY™ can cause cytokine release syndrome, including life-threatening or fatal reactions. In the clinical trial, CRS occurred in 76% of patients who received TALVEY™ at the recommended dosages, with Grade 1 CRS occurring in 57% of patients, Grade 2 in 17%, and Grade 3 in 1.5%. Recurrent CRS occurred in 30% of patients. CRS occurred in 33% of patients with step-up dose 3 in the biweekly dosing schedule (N=153). CRS occurred in 30% of patients with the first 0.4 mg/kg treatment dose and in 12% of patients treated with the first 0.8 mg/kg treatment dose. The CRS rate for both dosing schedules combined was less than 3% for each of the remaining doses in Cycle 1 and less than 3% cumulatively from Cycle 2 onward. The median time to onset of CRS was 27 (range: 0.1 to 167) hours from the last dose, and the median duration was 17 (range: 0 to 622) hours. Clinical signs and symptoms of CRS include but are not limited to pyrexia, hypotension, chills, hypoxia, headache, and tachycardia. Potentially life-threatening complications of CRS may include cardiac dysfunction, acute respiratory distress syndrome, neurologic toxicity, renal and/or hepatic failure, and disseminated intravascular coagulation (DIC).

Initiate therapy with step-up dosing and administer pre-treatment medications (corticosteroids, antihistamine, and antipyretics) prior to each dose of TALVEY™ in the step-up dosing schedule to reduce the risk of CRS. Monitor patients following administration accordingly. In patients who experience CRS, pre-treatment medications should be administered prior to the next TALVEY™ dose.

Counsel patients to seek medical attention should signs or symptoms of CRS occur. At the first sign of CRS, immediately evaluate patient for hospitalization and institute treatment with supportive care based on severity, and consider further management per current practice guidelines. Withhold TALVEY™ until CRS resolves or permanently discontinue based on severity.

Please read full [Prescribing Information](#), including **Boxed WARNING**, for TALVEY™.



6. Important Safety Information (continued)

Neurologic Toxicity including ICANS: TALVEY™ can cause serious or life-threatening neurologic toxicity, including immune effector cell-associated neurotoxicity syndrome (ICANS), including fatal reactions. In the clinical trial, neurologic toxicity occurred in 55% of patients who received the recommended dosages, with Grade 3 or 4 neurologic toxicity occurring in 6% of patients. The most frequent neurologic toxicities were headache (20%), encephalopathy (15%), sensory neuropathy (14%), and motor dysfunction (10%).

ICANS was reported in 9% of 265 patients where ICANS was collected and who received the recommended dosages. Recurrent ICANS occurred in 3% of patients. Most patients experienced ICANS following step-up dose 1 (3%), step-up dose 2 (3%), step-up dose 3 of the biweekly dosing schedule (1.8%), or the initial treatment dose of the weekly dosing schedule (2.6%) (N=156) or the biweekly dosing schedule (3.7%) (N=109). The median time to onset of ICANS was 2.5 (range: 1 to 16) days after the most recent dose with a median duration of 2 (range: 1 to 22) days. The onset of ICANS can be concurrent with CRS, following resolution of CRS, or in the absence of CRS. Clinical signs and symptoms of ICANS may include but are not limited to confusional state, depressed level of consciousness, disorientation, somnolence, lethargy, and bradyphrenia.

Monitor patients for signs and symptoms of neurologic toxicity during treatment. At the first sign of neurologic toxicity, including ICANS, immediately evaluate the patient and provide supportive care based on severity; withhold or permanently discontinue TALVEY™ based on severity and consider further management per current practice guidelines. [see Dosage and Administration (2.5)].

Due to the potential for neurologic toxicity, patients receiving TALVEY™ are at risk of depressed level of consciousness. Advise patients to refrain from driving or operating heavy or potentially dangerous machinery during the step-up dosing schedule and for 48 hours after completion of the step-up dosing schedule, and in the event of new onset of any neurological symptoms, until symptoms resolve.

TECVAYLI® and TALVEY™ REMS: TALVEY™ is available only through a restricted program under a REMS, called the TECVAYLI® and TALVEY™ REMS because of the risks of CRS and neurologic toxicity, including ICANS.

Further information about the TECVAYLI® and TALVEY™ REMS program is available at www.TEC-TALREMS.com or by telephone at 1-855-810-8064.

Oral Toxicity and Weight Loss: TALVEY™ can cause oral toxicities, including dysgeusia, dry mouth, dysphagia, and stomatitis. In the clinical trial, 80% of patients had oral toxicity, with Grade 3 occurring in 2.1% of patients who received the recommended dosages. The most frequent oral toxicities were dysgeusia (49%), dry mouth (34%), dysphagia (23%), and ageusia (18%). The median time to onset of oral toxicity was 15 (range: 1 to 634) days, and the median time to resolution to baseline was 43 (1 to 530) days. Oral toxicity did not resolve to baseline in 65% of patients.

TALVEY™ can cause weight loss. In the clinical trial, 62% of patients experienced weight loss of 5% or greater, regardless of having an oral toxicity, including 28% of patients with Grade 2 (10% or greater) weight loss and 2.7% of patients with Grade 3 (20% or greater) weight loss. The median time to onset of Grade 2 or higher weight loss was 67 (range: 6 to 407) days, and the median time to resolution was 50 (range: 1 to 403) days. Weight loss did not resolve in 57% of patients who reported weight loss.

Monitor patients for signs and symptoms of oral toxicity. Counsel patients to seek medical attention should signs or symptoms of oral toxicity occur and provide supportive care as per current clinical practice, including consultation with a nutritionist. Monitor weight regularly during therapy. Evaluate clinically significant weight loss further. Withhold TALVEY™ or permanently discontinue based on severity.

Please read full [Prescribing Information](#), including **Boxed WARNING**, for TALVEY™.



6. Important Safety Information (continued)

Infections: TALVEY™ can cause infections, including life-threatening or fatal infections. Serious infections occurred in 16% of patients, with fatal infections in 1.5% of patients. Grade 3 or 4 infections occurred in 17% of patients. The most common serious infections reported were bacterial infection (8%), which included sepsis and COVID-19 (2.7%).

Monitor patients for signs and symptoms of infection prior to and during treatment with TALVEY™ and treat appropriately. Administer prophylactic antimicrobials according to local guidelines. Withhold or permanently discontinue TALVEY™ as recommended, based on severity.

Cytopenias: TALVEY™ can cause cytopenias, including neutropenia and thrombocytopenia. In the clinical trial, Grade 3 or 4 decreased neutrophils occurred in 35% of patients, and Grade 3 or 4 decreased platelets occurred in 22% of patients who received TALVEY™. The median time to onset for Grade 3 or 4 neutropenia was 22 (range: 1 to 312) days, and the median time to resolution to Grade 2 or lower was 8 (range: 1 to 79) days. The median time to onset for Grade 3 or 4 thrombocytopenia was 12 (range: 2 to 183) days, and the median time to resolution to Grade 2 or lower was 10 (range: 1 to 64) days. Monitor complete blood counts during treatment and withhold TALVEY™ as recommended, based on severity.

Skin Toxicity: TALVEY™ can cause serious skin reactions, including rash, maculo-papular rash, erythema, and erythematous rash. In the clinical trial, skin reactions occurred in 62% of patients, with grade 3 skin reactions in 0.3%. The median time to onset was 25 (range: 1 to 630) days. The median time to improvement to grade 1 or less was 33 days.

Monitor for skin toxicity, including rash progression. Consider early intervention and treatment to manage skin toxicity. In the clinical trial, supportive care included topical steroids (15%). Oral steroid tapers (4.4%) were typically administered for Grade 3 skin reactions. Withhold or permanently discontinue TALVEY™, based on severity.

Hepatotoxicity: TALVEY™ can cause hepatotoxicity. Elevated ALT occurred in 33% of patients, with grade 3 or 4 ALT elevation occurring in 2.7%; elevated AST occurred in 31% of patients, with grade 3 or 4 AST elevation occurring in 3.3%. Grade 3 or 4 elevations of total bilirubin occurred in 0.3% of patients. Liver enzyme elevation can occur with or without concurrent CRS.

Monitor liver enzymes and bilirubin at baseline and during treatment as clinically indicated. Withhold TALVEY™ or consider permanent discontinuation of TALVEY™, based on severity [see Dosage and Administration (2.5)].

Embryo-Fetal Toxicity: Based on its mechanism of action, TALVEY™ may cause fetal harm when administered to a pregnant woman. Advise pregnant women of the potential risk to the fetus. Advise females of reproductive potential to use effective contraception during treatment with TALVEY™ and for 3 months after the last dose.

Adverse Reactions: The most common adverse reactions ($\geq 20\%$) are pyrexia, CRS, dysgeusia, nail disorder, musculoskeletal pain, skin disorder, rash, fatigue, weight decreased, dry mouth, xerosis, dysphagia, upper respiratory tract infection, diarrhea, hypotension, and headache.

The most common Grade 3 or 4 laboratory abnormalities ($\geq 30\%$) are lymphocyte count decreased, neutrophil count decreased, white blood cell decreased, and hemoglobin decreased.

cp-394174v2

Please read full [Prescribing Information](#), including **Boxed WARNING**, for TALVEY™.



[illegible]

TALVEY™
(talquetamab-tgvs) Injection for
subcutaneous use
2 mg/mL and 40 mg/mL

Please read full Important Safety Information on [pages 21-23](#) and full [Prescribing Information](#), including Boxed WARNING, for TALVEY™.

Reference: TALVEY™ [Prescribing Information]. Horsham, PA: Janssen Biotech, Inc.

iKnowMed® is a registered trademark of McKesson.
For more information, visit www.TALVEYhcp.com.

