

Ongoing



or



TECVAYLI® and TALVEY® are two distinct bispecific antibodies with differences in dosing and administration, safety, and use.

This reference guide provides helpful information for ongoing treatment with TECVAYLI® or TALVEY®. This is not intended to be a comprehensive list, and individual practices must follow their own required procedures.

This checklist is intended for institutions who provide ongoing treatment for TECVAYLI® or TALVEY®.

Providers may utilize this checklist for their practice while either referring patients to an initiating treatment center or providing ongoing treatment.



Referring Patient to Initiating Treatment Center

- ☐ Determine the appropriate initiating treatment center for your patient based on Risk Evaluation and Mitigation Strategy (REMS) certification, geographical distance, and insurance coverage
 - ☐ Please visit [TECVAYLI.com](https://www.tecvayli.com) or [TALVEY.com](https://www.talvey.com) to locate a REMS-certified treatment center with the TECVAYLI® Treatment Locator Tool/TALVEY® Treatment Locator Tool.
- ☐ Align on patient treatment care plan and coordinate patient transfer with the initiating treatment center
 - ☐ Establish effective communication methods with initiating treatment provider(s) and/or care team
 - ☐ Forward any important discharge information, treatment notes, and records to initiating treatment site
 - ☐ Outline and discuss patient return for ongoing treatment
- ☐ Encourage the patient or their caregiver to engage the initiating treatment center for bispecific education, including outlining the step-up dosing schedule and administrative logistics



Preparing for Ongoing Treatment

- ☐ Ensure that treating prescribers and pharmacists receive REMS certification
- ☐ To ensure a successful transfer of the patient, communicate with initiating treatment site on status of the treatment plan and timing of transfer, and request all necessary transfer information (treatment notes and patient records)
- ☐ Perform benefits investigation (eg, J-Code, New Technology Add-On Payments [NTAP]) and obtain new prior authorization, if required
- ☐ Establish adverse reaction (AR) protocols and train the corresponding staff on Cytokine Release Syndrome (CRS), Neurologic Toxicity, including Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS), and infections, as well as TALVEY®-specific AR management incidence and protocols (eg, Nurse, Advanced Practice Provider, Pharmacist, Dermatologist, Social Worker, Nutritionist, Emergency Department, Infectious Disease)
- ☐ Dosing ordering information:
 - ☐ **TECVAYLI®:** Order 153 mg/1.7 mL vial size for ongoing treatment dose; use Table 9 in the full Prescribing Information to determine number of vials required based on patient's actual body weight
 - ☐ **TALVEY®:** Order 40 mg/mL vial size for ongoing biweekly or weekly treatment dose; use Tables 11 and 12 in the full Prescribing Information to determine number of vials required based on patient's actual body weight
- ☐ Ensure supportive therapies are available for CRS and Neurologic Toxicity, including ICANS
 - ☐ Please see the *TECVAYLI® Dosing and AR Management Guide*, the *TALVEY® Treatment Management Guide* or the *TECVAYLI® and TALVEY® Adverse Reaction Management Guide* for reference.
 - ☐ Consider asking your local Johnson & Johnson representative and/or Oncology Clinical Educator for more information on any AR management
- ☐ Confirm the patient has received and is carrying the Patient Wallet Card for either TECVAYLI® or TALVEY® as prescribed
- ☐ Consider connecting with your local Johnson & Johnson representative to receive any necessary resources or if there are any questions



Patient/Caregiver Education and Logistics for Ongoing Treatment

- ☐ Discuss the patient's step-up dosing experience and ensure the patient and caregiver understand the ongoing treatment plan for TECVAYLI® or TALVEY®
- ☐ Please see section 17 in the respective product's full Prescribing Information for patient counseling information. Also, please provide and advise the patient to read the respective product's Medication Guide
- ☐ Ensure the patient has scheduled their next treatment doses, while confirming support during treatment, including caregiver support, travel, and transportation to/from appointments
- ☐ Discuss any concerns with the patient and caregiver about ARs and recap the treatment plan if the patient experiences any symptoms of ARs
 - ☐ Provide the phone number of the treating physician and where to go if the patient experiences any ARs
 - ☐ Please see section 17 in the respective product's full Prescribing Information for patient counseling information
 - ☐ Please work with your local Johnson & Johnson representative and refer the patient and caregiver to any key resources that would be helpful

Ongoing

TECVAYLI®
(teclistamab-cqyv) Injection for
subcutaneous use
10 mg/mL and 90 mg/mL

or

TALVEY®
(talquetamab-tgvs) Injection for
subcutaneous use
2 mg/mL and 40 mg/mL



Helpful Resources for Your Reference



TECVAYLI® Prescribing Information



TALVEY® Prescribing Information



**REMS Fact Sheet and
Cover Letter**



TECVAYLI® Medication Guide



TALVEY® Medication Guide



**Janssen CarePath
Resource Guide**



**TECVAYLI® Dosing and AR
Management Guide**



**TALVEY® Treatment
Management Guide**



TECVAYLI® Patient Brochure



TALVEY® Patient Brochure



TECVAYLI® Getting Started Guide



My Journey With TALVEY®



TECVAYLI® Caregiver Brochure



TALVEY® Caregiver Brochure



*Scan the QR code to connect with a local Johnson & Johnson representative
and learn more about these resources*